



Caledonian Structures of the Lake District (CAL-STR-LD)

Block Description

Visit <https://jncc.gov.uk/gcr-site-list>, for more information on GCR blocks and sites
For Precambrian and Structural Geology GCR block descriptions and GCR site lists,
visit <https://jncc.gov.uk/gcr-blocks-precambrian-and-structural-geology>

Introduction

The GCR sites selected for this GCR Block represent a major episode of Earth history that occurred as a result of the 'Caledonian Orogeny', lasting from about 500 million years ago (Ma) to around 360 Ma, which was a period of mountain building and continental collision that began in the Ordovician Period (which lasted from 495 to 440 Ma) and continued through Silurian (440–417 Ma) and Devonian (417–354 Ma) time, as represented by geological features present in the Lake District; geological structures south of the Southern Upland Fault produced elsewhere by the same events are covered by other Blocks: the Caledonian Structures of the Southern Uplands and Caledonian Structures of Wales, for both geographical and geological reasons. See the GCR Blocks **Caledonian Structures of the Southern Uplands (CAL-STR-SU)** and **Caledonian Structures of Wales (CAL-STR-WL)**. Igneous rocks that formed as a result of the Caledonian Orogeny are encompassed by the **Caledonian igneous (CAL-IGN)** GCR Block

There is at present no agreed definition of the term 'Caledonian', but is taken here to include all of the convergent tectonic and magmatic events arising from the closure of the 'proto-Atlantic' Iapetus Ocean in which many of the rocks of Late Proterozoic and early Palaeozoic age had been deposited. It therefore encompasses subduction beneath the continental margins; the accretion or obduction of oceanic crust and island-arc material onto these margins; and ultimate collision of the continents, uplift and development of extensional molasse basins. Within this broad orogenic framework many separate 'events' are identified, such as the 'Grampian Event' and the 'Acadian Event'.

For details of stratigraphy related to the Caledonian Orogeny see GCR Blocks for Ordovician, Silurian and Devonian Stratigraphy: **Arenig - Llanvirn (ARE-LLV)**; **Arenig - Tremadoc (ARE-TRE)**; **Caradoc-Ashgill (CAR-ASH)**; **Llandeilo (LDO)**; **Llandovery (LDY)**; **Ludlow (LUD)**; **Marine Devonian (MAR-DEV)**; **Non-Marine Devonian (NMAR-DEV)**.

Tectonic setting and evolution

The Iapetus Ocean was created in Late Proterozoic time by the rifting and pulling apart of a large supercontinent known as 'Rodinia'. The opening started sometime around 650 million years (Ma) ago and by the beginning of Ordovician time, at 510 Ma, the ocean was at its widest development of possibly up to 5000 km across. On one side of the ocean lay the supercontinent of Laurentia, which is represented today largely by the Precambrian basement rocks of North America, Greenland, the north of Ireland and the Scottish Highlands. On the opposite side lay the supercontinent of Gondwana, consisting of the basements of South America, Africa, India, Australia, East Antarctica and Western Europe (including south Ireland, England and Wales). A separate continent, 'Baltica' (the basement of Scandinavia and Russia), was separated from Gondwana by an arm of the Iapetus Ocean, known as the 'Tornquist Sea'. The wide separation is supported by palaeontological data, which shows distinctly different faunal assemblages in the Lower Palaeozoic rocks of each continent and by palaeomagnetic interpretations.

The continental plates of Laurentia, Gondwana and Baltica started to converge during the early part of the Ordovician Period, initiating new tectonic and magmatic processes which marked the start of the Caledonian Orogeny. The Iapetus oceanic crust was consumed in subduction zones beneath oceanic island arcs and beneath the continental margins. Magma was being created by the melting of mantle and oceanic crustal material within and above the subduction zones and by melting within the thickened continental crust.

The generally accepted sequence and timing of events as the three plates converged is as follows.

- Closure of the Tornquist Sea between Eastern Avalonia and Baltica, followed by strike-slip movement along the Tornquist Suture.

- Anticlockwise rotation of Baltica, followed by convergence with Laurentia, with subduction beneath the 'Scottish' sector of the Laurentian margin and closure during early to mid-Silurian times.
- Oblique convergence of Eastern Avalonia with Laurentia, with subduction beneath the Laurentian margin, resulting in closure by the early Silurian in the 'Irish' sector and later, mid-Silurian closure in the 'Scottish' sector. The junction between the two fused plates passes through the Solway Firth in Britain and is known as the Iapetus Suture.
- Protracted continent–continent collision between Laurentia and Eastern Avalonia plus Baltica, with underthrusting beneath part of the Laurentian margin (?mid-Silurian to Mid-Devonian).
- Separation of a further microcontinent, Armorica, from the margin of Gondwana, which then collided with Eastern Avalonia during Early Devonian times (the Acadian Event).
- Sinistral re-alignment of terrane boundaries (?mid-Silurian to Mid-Devonian).

The Caledonian Orogenic Belt

The Caledonian–Appalachian Orogen can be traced (pre-Atlantic drift), for some 7500 km south-west to north-east, from south-eastern USA through the British Isles to Scandinavia, Greenland, and Ny Friesland. It is generally accepted that sedimentation and igneous activity took place at, or near, the margins of an ocean (the Iapetus) that separated the Laurentian and Gondwanaland plates, over a period from the Precambrian through the early Palaeozoic. From studies of fauna, sedimentary history, igneous activity, structural and metamorphic evolution, and palaeomagnetism on its two sides, it is considered that deformation of sediments and volcanic rocks, resulting from the episodic closure of the Iapetus Ocean, took place through the early Palaeozoic to culminate in continental collision during the early part of the Devonian Period.

The British Caledonides–Lake District

On the south-eastern margin, the argument for SE-directed subduction rests largely on the presence of arc volcanism in the Lake District during the Llandeilo or Caradoc epochs. Structural arguments have focused on the nature and significance of the pre- and post-Borrowdale Volcanic Group unconformities, but particular features that could be related directly to subduction have not been identified. Folding in the Skiddaw Group has been re-interpreted as being the product of slumping, and this may prove important with regard to the arguments for the timing of subduction and the topography of the margin. Apart from tilting and block faulting due to volcanotectonic activity, the dominant deformation in the Lake District is regarded as essentially a single event that resulted in folding, cleavage, and greenschist metamorphism during the early Devonian.